

# T-wave inversions during conduction system pacing: A marker of more physiological ventricular activation

Karol Čurila<sup>1</sup>, Pavel Jurak<sup>2</sup>, Uyen Chau Nguyen<sup>3</sup>

## Related article

by Geng et al.

<sup>1</sup>Cardiocenter, 3<sup>rd</sup> Faculty of Medicine, Charles University and University Hospital Kralovske Vinohrady, Prague, Czech Republic

<sup>2</sup>Institute of Scientific Instruments, the Czech Academy of Sciences, Brno, Czech Republic

<sup>3</sup>Department of Physiology and Cardiology, Maastricht University Medical Center, Maastricht, the Netherlands

### Correspondence to:

Karol Čurila, MD, PhD,  
Cardiocenter, 3<sup>rd</sup> Faculty of  
Medicine, Charles University and  
University Hospital Kralovske  
Vinohrady,  
Srobarova 50, 100 34, Praha 10,  
Czech Republic,  
phone: +42 0 777 869 171,  
e-mail: karol.curila@fnkv.cz

Copyright by the Author(s), 2022

DOI: 10.33963/KPa2022.

### Received:

October 11, 2022

### Accepted:

October 11, 2022

### Early publication date:

October 31, 2022


Permanent myocardial pacing can preserve adequate heart rates and improve symptoms and mortality in patients with bradycardia [1]. Conventional right ventricular (RV) pacing is far from the optimal treatment since up to 20% of patients experience a reduction in the left ventricular (LV) ejection fraction, which can lead to heart failure (HF) [2]. This was the main incentive for developing 'conduction system pacing (CSP)' techniques that target (directly or indirectly) the capture of conduction tissue, initiating more physiological ventricular activation. Although His bundle pacing (HBP) leads to the best ventricular synchrony [3], proper positioning is complicated, requires high pacing thresholds, and is associated with lower sensing values. For these reasons, left bundle branch area pacing (LBBAP), where the lead tip is deployed in the left subendocardial septal area, is currently preferred over HBP.

While ventricular activation is reasonably well understood [4, 5], little is known about the repolarization sequence during LBBAP. In a sizeable group of patients, Geng et al. [6] investigated the effect of LBBAP on occurrence and characteristics of T-wave inversions (TWIs). They showed that TWIs frequently (in 66% of cases) occurred one day after initiating LBBAP. TWIs appeared more frequently in patients with bundle branch blocks, and the main TWI predictor was QRS duration  $\geq 120$  ms. TWIs were unrelated to myocardial ischemia and, in most patients (88%), partially

or entirely disappeared during a median follow-up of 10 days.

It is 40 years since Rosenbaum described the occurrence of transient TWIs in animal experiments [7]. He showed that temporary ventricular pacing could lead to changes in T-wave vectors and polarities that persisted after cessation of pacing and restoration of physiological ventricular activation. Interestingly, T-wave inversions were observed in the same leads in which the polarities of QRS complexes had been changed during pacing. Rosenbaum coined the term "cardiac memory" to note this phenomenon. This phenomenon has been observed in patients after cessation of RV pacing, successful ablation of accessory pathways associated with intermittent LBBB and ventricular tachyarrhythmias [8].

In a normal heart, the last activated regions of the ventricular wall generally have shorter action potential durations (APD) compared to the ones activated earlier. Myocytes that depolarize last will therefore repolarize first [9]. However, when the ventricular activation sequence suddenly changes, the distribution of APD does not immediately adapt to this change. Consequently, ventricular wall regions that were previously activated late (with shorter APD) may become activated sooner, creating a situation where myocytes that depolarize first also repolarize first. A similar situation was reported in patients with HF and wide QRS complexes [9].



**Figure 1.** Visual representation of regional T-wave inversion distribution in LBBB (left) and RBBB (right) patients. Data presented in Table 3 in the study by Geng et al. [6] was used as input. The regional TWI percentages were computed by averaging the percentages belonging to leads V1 and V2 (septum), V2 and V3 (anterior), V5, V6, I, and aVL (lateral), II, III, and aVF (inferior)

Abbreviations: LBBB, left bundle branch block; RBBB, right bundle branch block; TWI, T-wave inversion

This property appears to explain the observation by Geng et al. that TWI locations depend on the type of the bundle branch block; i.e., in LBBB patients, they occurred more frequently in leads V1–V4, II, III, and aVF, but in RBBB patients, leads I and aVL were affected more often. Figure 1 visualizes the incidence of TWIs per ECG-lead using data from Geng et al. (Table 3) [6] and assumes the following lead associations: leads V1 and 2 with the septum, V2 and V3 the anterior, V5, V6, I, and aV — the lateral wall, and II, III, and aVF — the inferior wall. In this representation (Figure 1), it becomes clear that TWIs in LBBB patients occurred most often in regions in which the sequence of ventricular activation and polarities of QRS complexes changed significantly during LBBAP. It would be interesting to determine if TWIs distribution in RBBB patients was associated with fascicular hemiblocks, which lead to less physiological LV activation. That was, however, not analyzed by the authors. Another reason may be including the patients with heart failure, which may lead to the inclusion of altered activation-repolarization relationships even in the absence of conduction disturbances [9]. Few studies have investigated repolarization changes following biventricular pacing in HF patients. Dispersion of repolarization appears to have a time-dependent character, with a high amount of dispersion observed within one month after implantation and decreasing over time [10].

Interestingly, computer simulations have demonstrated that acute biventricular pacing reduces LV repolarization dispersion on a regional level while increasing RV repolarization dispersion, leading to a higher degree of interventricular repolarization dispersion. During chronic biventricular pacing, however, an adaptation of APDs occurs, leading to a reduction in repolarization dispersion [10]. These processes are compatible with physiological adaptations to minimize dispersion of repolarization, so the disappearance of TWIs appears to be a physiological process.

Although inverted T-waves concern cardiologists, it is unclear to what extent TWIs related to cardiac memory are associated with an elevated risk for arrhythmias. Most clinical studies investigating the efficacy of conventional

biventricular pacing did not find an association with ventricular tachyarrhythmias [11–13]. However, multiple individual cases describing the occurrence of electrical storm shortly after CRT implantation raised concerns about the pro-arrhythmic effect of LV epicardial pacing [14]. While Geng et al. [6] did not investigate T-wave changes during conventional biventricular pacing, Gupta et al. [15] recently demonstrated that CSP (both HBP and LBBAP) was associated with reduced repolarization heterogeneity (defined as Tpeak-Tend on a surface ECG) and greater cardiac memory resolution compared to conventional biventricular pacing. Additional studies comparing the temporal evolution of repolarization changes during biventricular vs. conduction system pacing are certainly warranted. More mechanistic insights can be obtained using invasive or non-invasive electro-anatomic mapping techniques.

### Article information

**Acknowledgments:** The authors thank Prof. Frits W. Prinzen for his scientific contribution to this editorial and Dr. R. Meiburg for his assistance in preparing Figure 1.

**Conflict of interest:** None declared.

**Funding:** This work was supported by the National Institute for Research of Metabolic and Cardiovascular Diseases project (Programme EXCELES, ID Project No. LX22NPO5104) — Funded by the European Union — Next Generation EU and a personal grant from the Dutch Heart Foundation (2021T016).

**Open access:** This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at [kardiologiapolska@ptkardio.pl](mailto:kardiologiapolska@ptkardio.pl).

### REFERENCES

- Glikson M, Nielsen JC, Kronborg MB, et al. 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy. *Europace*. 2022; 24(1): 71–164, doi: [10.1093/europace/euab232](https://doi.org/10.1093/europace/euab232), indexed in PubMed: 34455427.
- Kiehl EL, Makki T, Kumar R, et al. Incidence and predictors of right ventricular pacing-induced cardiomyopathy in patients with complete atrioventricular block and preserved left ventricular systolic function. *Heart Rhythm*. 2016; 13(12): 2272–2278, doi: [10.1016/j.hrthm.2016.09.027](https://doi.org/10.1016/j.hrthm.2016.09.027), indexed in PubMed: 27855853.
- Curila K, Jurak P, Halamek J, et al. Ventricular activation pattern assessment during right ventricular pacing; ultra-high-frequency ECG study. *J Cardiovasc Electrophysiol*. 2021; 32(5): 1385–1394, doi: [10.1111/jce.14985](https://doi.org/10.1111/jce.14985), indexed in PubMed: 33682277.
- Curila K, Jurak P, Jastrzebski M, et al. Left bundle branch pacing compared to left ventricular septal myocardial pacing increases interventricular dyssynchrony but accelerates left ventricular lateral wall depolarization. *Heart Rhythm*. 2021; 18(8): 1281–1289, doi: [10.1016/j.hrthm.2021.04.025](https://doi.org/10.1016/j.hrthm.2021.04.025), indexed in PubMed: 33930549.
- Curila K, Jurak P, Vernooij K, et al. Left Ventricular Myocardial Septal Pacing in Close Proximity to LBB Does Not Prolong the Duration of the Left Ventricular Lateral Wall Depolarization Compared to LBB Pacing. *Front Cardiovasc Med*. 2021; 8: 787414, doi: [10.3389/fcvm.2021.787414](https://doi.org/10.3389/fcvm.2021.787414), indexed in PubMed: 34950718.
- Geng J, Jiang Z, Zhang S, et al. Reversible T-wave inversions during left bundle branch area pacing. *Kardiol Pol*. 2022; 80(10): 1002–1009, doi: [10.33963/KP.a2022.0167](https://doi.org/10.33963/KP.a2022.0167), indexed in PubMed: 35836370.
- Rosenbaum M, Blanco H, Elizari M, et al. Electrotonic modulation of the T wave and cardiac memory. *Am J Cardiol* 1982; 50(2): 213–222, doi: [10.1016/0002-9149\(82\)90169-2](https://doi.org/10.1016/0002-9149(82)90169-2).

8. Jeyaraj D, Ashwath M, Rosenbaum DS. Pathophysiology and clinical implications of cardiac memory. *Pacing Clin Electrophysiol.* 2010; 33(3): 346–352, doi: [10.1111/j.1540-8159.2009.02630.x](https://doi.org/10.1111/j.1540-8159.2009.02630.x), indexed in Pubmed: 20025710.
9. Maffessanti F, Wanten J, Potse M, et al. The relation between local repolarization and T-wave morphology in heart failure patients. *Int J Cardiol.* 2017; 241: 270–276, doi: [10.1016/j.ijcard.2017.02.056](https://doi.org/10.1016/j.ijcard.2017.02.056), indexed in Pubmed: 28318665.
10. Verzaal NJ, van Deursen CJM, Pezzuto S, et al. Synchronization of repolarization after cardiac resynchronization therapy: A combined clinical and modeling study. *J Cardiovasc Electrophysiol.* 2022; 33(8): 1837–1846, doi: [10.1111/jce.15581](https://doi.org/10.1111/jce.15581), indexed in Pubmed: 35662306.
11. Roque C, Trevisi N, Silberbauer J, et al. Electrical storm induced by cardiac resynchronization therapy is determined by pacing on epicardial scar and can be successfully managed by catheter ablation. *Circ Arrhythm Electrophysiol.* 2014; 7(6): 1064–1069, doi: [10.1161/CIRCEP.114.001796](https://doi.org/10.1161/CIRCEP.114.001796), indexed in Pubmed: 25221332.
12. Cabanelas N, Oliveira M, Nogueira da Silva M, et al. The proarrhythmic effect of cardiac resynchronization therapy: an issue that should be borne in mind. *Rev Port Cardiol.* 2014; 33(5): 309.e1–309.e7, doi: [10.1016/j.repc.2014.01.011](https://doi.org/10.1016/j.repc.2014.01.011), indexed in Pubmed: 24931180.
13. Medina-Ravell VA, Lankipalli RS, Yan GX, et al. Effect of epicardial or biventricular pacing to prolong QT interval and increase transmural dispersion of repolarization: does resynchronization therapy pose a risk for patients predisposed to long QT or torsade de pointes? *Circulation.* 2003; 107(5): 740–746, doi: [10.1161/01.cir.0000048126.07819.37](https://doi.org/10.1161/01.cir.0000048126.07819.37), indexed in Pubmed: 12578878.
14. Gold MR, Linde C, Abraham WT, et al. The impact of cardiac resynchronization therapy on the incidence of ventricular arrhythmias in mild heart failure. *Heart Rhythm.* 2011; 8(5): 679–684, doi: [10.1016/j.hrthm.2010.12.031](https://doi.org/10.1016/j.hrthm.2010.12.031), indexed in Pubmed: 21185401.
15. Gupta A, Pavri BB. Conduction system pacing versus biventricular pacing: Reduced repolarization heterogeneity in addition to improved depolarization. *J Cardiovasc Electrophysiol.* 2022; 33(2): 287–295, doi: [10.1111/jce.15329](https://doi.org/10.1111/jce.15329), indexed in Pubmed: 34911154.